

FLUDEOXYGLUCOSE F 18 - fludeoxyglucose f-18 injection
The University of Utah DBA Cyclotron Radiochemistry Lab / Huntsman Cancer Institute

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use Fludeoxyglucose F18 Injection, USP safely and effectively. See full prescribing information for Fludeoxyglucose F18 Injection, USP.

Fludeoxyglucose F18 Injection, USP

For intravenous use

Initial U.S. Approval: 2005

----- **INDICATIONS AND USAGE** -----

Fludeoxyglucose F18 Injection, USP is indicated for positron emission tomography (PET) imaging in the following settings: (1)

- Oncology: For assessment of abnormal glucose metabolism to assist in the evaluation of malignancy in patients with known or suspected abnormalities found by other testing modalities, or in patients with an existing diagnosis of cancer. (1)
- Cardiology: For the identification of left ventricular myocardium with residual glucose metabolism and reversible loss of systolic function in patients with coronary artery disease and left ventricular dysfunction, when used together with myocardial perfusion imaging. (1)
- Neurology: For the identification of regions of abnormal glucose metabolism associated with foci of epileptic seizures (1). (1)

----- **DOSAGE AND ADMINISTRATION** -----

Fludeoxyglucose F18 Injection emits radiation. Use procedures to minimize radiation exposure. Screen for blood glucose abnormalities. (2)

• In the oncology and neurology settings, instruct patients to fast for 4 – 6 hours prior to the drug's injection. Consider medical therapy and laboratory testing to assure at least two days of normoglycemia prior to the drug's administration (5.2). (2)

• In the cardiology setting, administration of glucose-containing food or liquids (e.g., 50 – 75 grams) prior to the drug's injection facilitates localization of cardiac ischemia (2.3). (2)

Aseptically withdraw Fludeoxyglucose F18 Injection from its container and administer by intravenous injection (2). The recommended dose: (2)

- for adults is 5 – 10 mCi (185 – 370 MBq), in all indicated clinical settings (2.1). (2)
- for pediatric patients is 2.6 mCi in the neurology setting (2.2). Initiate imaging within 40 minutes following drug injection; acquire static emission images 30 – 100 minutes from time of injection (2). (2)

----- **DOSAGE FORMS AND STRENGTHS** -----

Multiple-dose glass vial containing 0.74 – 11.1 GBq (20 – 300 mCi/mL) of Fludeoxyglucose F18 Injection and 4.5 mg of sodium chloride in citrate buffer (approximately 23 mL or 35 mL volume), for intravenous administration (3). (3)

----- **CONTRAINDICATIONS** -----

None (4)

----- **WARNINGS AND PRECAUTIONS** -----

- Radiation risks: use smallest dose necessary for imaging (5.1). (5)
- Blood glucose abnormalities: may cause suboptimal imaging (5.2). (5)

----- **ADVERSE REACTIONS** -----

Hypersensitivity reactions have occurred; have emergency resuscitation equipment and personnel immediately available (6). (6)

(6)

(6)

To report SUSPECTED ADVERSE REACTIONS, contact The University of Utah DBA Cyclotron Radiochemistry Lab Huntsman Cancer Institute (6)

at 801-581-8745 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. (6)

(6)

----- **USE IN SPECIFIC POPULATIONS** -----

- Pregnancy Category C: No human or animal data. Consider alternative diagnostics; use only if clearly needed (8.1). (8)
- Nursing mothers: Use alternatives to breast feeding (e.g., stored breast milk or infant formula) for at least 10 half-lives of radioactive decay, if Fludeoxyglucose F 18 Injection is administered to a woman who is breastfeeding(8.3). (8)
- Pediatric Use: Safety and effectiveness in pediatric patients have not been established in the oncology and cardiology settings (8.4). (8)

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 6/2015

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

- 1.1 Oncology**
- 1.2 Cardiology**
- 1.3 Neurology**

2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dose for Adults**
- 2.2 Recommended Dose for Pediatric Patients**
- 2.3 Patient Preparation**
- 2.4 Radiation Dosimetry**
- 2.5 Radiation Safety – Drug Handling**
- 2.6 Drug Preparation and Administration**
- 2.7 Imaging Guidelines**

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Radiation Risks**
- 5.2 Blood Glucose Abnormalities**

6 ADVERSE REACTIONS

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy**
- 8.3 Nursing Mothers**
- 8.4 Pediatric Use**

11 DESCRIPTION

- 11.1 Chemical Characteristics**
- 11.2 Physical Characteristics**

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action**
- 12.2 Pharmacodynamics**
- 12.3 Pharmacokinetics**

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

14 CLINICAL STUDIES

- 14.1 Oncology**
- 14.2 Cardiology**
- 14.3 Neurology**

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND DRUG HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

Fludeoxyglucose F-18 Injection, USP is indicated for positron emission tomography (PET) imaging in the following settings:

1.1 Oncology

For assessment of abnormal glucose metabolism to assist in the evaluation of malignancy in patients with known or suspected abnormalities found by other testing modalities, or in patients with an existing diagnosis of cancer.

1.2 Cardiology

For the identification of left ventricular myocardium with residual glucose metabolism and reversible loss of systolic function in patients with coronary artery disease and left ventricular dysfunction, when used together with myocardial perfusion imaging.

1.3 Neurology

For the identification of regions of abnormal glucose metabolism associated with foci of epileptic seizures.

2 DOSAGE AND ADMINISTRATION

Fludeoxyglucose F-18 Injection emits radiation. Use procedures to minimize radiation exposure. Calculate the final dose from the end of synthesis (EOS) time using proper radioactive decay factors. Assay the final dose in a properly calibrated dose calibrator before administration to the patient [see Description(11.2)].

2.1 Recommended Dose for Adults

Within the oncology, cardiology and neurology settings, the recommended dose for adults is 5 – 10 mCi (185 – 370 MBq) as an intravenous injection.

2.2 Recommended Dose for Pediatric Patients

Within the neurology setting, the recommended dose for pediatric patients is 2.6 mCi, as an intravenous injection. The optimal dose adjustment on the basis of body size or weight has not been determined [see Use in Special Populations (8.4)].

2.3 Patient Preparation

- To minimize the radiation absorbed dose to the bladder, encourage adequate hydration. Encourage the patient to drink water or other fluids (as tolerated) in the 4 hours before their PET study.
- Encourage the patient to void as soon as the imaging study is completed and as often as possible thereafter for at least one hour.
- Screen patients for clinically significant blood glucose abnormalities by obtaining a history and/or laboratory tests [see Warnings and Precautions (5.2)]. Prior to Fludeoxyglucose F-18 PET imaging in the oncology and neurology settings, instruct patient to fast for 4 – 6 hours prior to the drug's injection.
- In the cardiology setting, administration of glucose-containing food or liquids (e.g., 50 – 75 grams) prior to Fludeoxyglucose F-18 Injection facilitates localization of cardiac ischemia.

2.4 Radiation Dosimetry

The estimated human absorbed radiation doses (rem/mCi) to a newborn (3.4 kg), 1-year old (9.8 kg), 5-year old (19 kg), 10-year old (32 kg), 15-year old (57 kg), and adult (70 kg) from intravenous administration of Fludeoxyglucose F-18 Injection are shown in Table 1. These estimates were calculated based on human² data and using the data published by the International Commission on Radiological Protection⁴ for Fludeoxyglucose¹⁸F. The dosimetry data show that there are slight

variations in absorbed radiation dose for various organs in each of the age groups. These dissimilarities in absorbed radiation dose are due to developmental age variations (e.g., organ size, location, and overall metabolic rate for each age group). The identified critical organs (in descending order) across all age groups evaluated are the urinary bladder, heart, pancreas, spleen, and lungs.

Table 1. Estimated Absorbed Radiation Doses (rem/mCi) After Intravenous Administration of Fludeoxyglucose F 18 Injection*

Organ	Newborn (3.4 kg)	1-year old (9.8 kg)	5-year old (19 kg)	10-year old (32 kg)	15-year old (57 kg)	Adult (70 kg)
Bladder wall†	4.3	1.7	0.93	0.60	0.40	0.32
Heart wall	2.4	1.2	0.70	0.44	0.29	0.22
Pancreas	2.2	0.68	0.33	0.25	0.13	0.096
Spleen	2.2	0.84	0.46	0.29	0.19	0.14
Lungs	0.96	0.38	0.20	0.13	0.092	0.064
Kidneys	0.81	0.34	0.19	0.13	0.089	0.074
Ovaries	0.80	0.8	0.19	0.11	0.058	0.053
Uterus	0.79	0.35	0.19	0.12	0.076	0.062
LLI wall‡	0.69	0.28	0.15	0.097	0.060	0.051
Liver	0.69	0.31	0.17	0.11	0.076	0.058
Gallbladder wall	0.69	0.26	0.14	0.093	0.059	0.049
Small intestine	0.68	0.29	0.15	0.096	0.060	0.047
ULI wall□	0.67	0.27	0.15	0.090	0.057	0.046
Stomach wall	0.65	0.27	0.14	0.089	0.057	0.047
Adrenals	0.65	0.28	0.15	0.095	0.061	0.048
Testes	0.64	0.27	0.14	0.085	0.052	0.041
Red marrow	0.62	0.26	0.14	0.089	0.057	0.047
Thymus	0.61	0.26	0.14	0.086	0.056	0.044
Thyroid	0.61	0.26	0.13	0.080	0.049	0.039
Muscle	0.58	0.25	0.13	0.078	0.049	0.039
Bone surface	0.57	0.24	0.12	0.079	0.052	0.041
Breast	0.54	0.22	0.11	0.068	0.043	0.034
Skin	0.49	0.20	0.10	0.060	0.037	0.030
Brain	0.29	0.13	0.09	0.078	0.072	0.070
Other tissues	0.59	0.25	0.13	0.083	0.052	0.042

* MIRDOSE 2 software was used to calculate the radiation absorbed dose. Assumptions on the biodistribution based on data from Gallagher et al.¹ and Jones et al.²

† The dynamic bladder model with a uniform voiding frequency of 1.5 hours was used.

‡ LLI = lower large intestine;

□ ULI = upper large intestine

2.5 Radiation Safety – Drug Handling

- Use waterproof gloves, effective radiation shielding, and appropriate safety measures when handling Fludeoxyglucose F-18 Injection to avoid unnecessary radiation exposure to the patient, occupational workers, clinical personnel and other persons.
- Radiopharmaceuticals should be used by or under the control of physicians who are qualified by specific training and experience in the safe use and handling of radionuclides, and whose experience and training have been approved by the appropriate governmental agency authorized to

license the use of radionuclides.

- Calculate the final dose from the end of synthesis (EOS) time using proper radioactive decay factors. Assay the final dose in a properly calibrated dose calibrator before administration to the patient [see Description (11.2)].
- The dose of Fludeoxyglucose F-18 used in a given patient should be minimized consistent with the objectives of the procedure, and the nature of the radiation detection devices employed.

2.6 Drug Preparation and Administration

- Calculate the necessary volume to administer based on calibration time and dose.
- Aseptically withdraw Fludeoxyglucose F-18 Injection from its container.
- Inspect Fludeoxyglucose F-18 Injection visually for particulate matter and discoloration before administration, whenever solution and container permit.
- Do not administer the drug if it contains particulate matter or discoloration; dispose of these unacceptable or unused preparations in a safe manner, in compliance with applicable regulations.
- Use Fludeoxyglucose F-18 Injection within 12 hours from the EOS

2.7 Imaging Guidelines

- Initiate imaging within 40 minutes following Fludeoxyglucose F-18 Injection administration.
- Acquire static emission images 30 – 100 minutes from the time of injection.

3 DOSAGE FORMS AND STRENGTHS

Multiple-dose glass vial containing 0.74 - 11.1GBq (20 - 300 mCi/mL) of Fludeoxyglucose F-18 Injection and 4.5 mg of sodium chloride in citrate buffer (approximately 23 mL or 35 mL volume), for intravenous administration.

4 CONTRAINDICATIONS

None

5 WARNINGS AND PRECAUTIONS

5.1 Radiation Risks

Radiation-emitting products, including Fludeoxyglucose F-18 Injection, may increase the risk for cancer, especially in pediatric patients. Use the smallest dose necessary for imaging and ensure safe handling to protect the patient and health care worker [see Dosage and Administration(2.5)].

5.2 Blood Glucose Abnormalities

In the oncology and neurology setting, suboptimal imaging may occur in patients with inadequately regulated blood glucose levels. In these patients, consider medical therapy and laboratory testing to assure at least two days of normoglycemia prior to Fludeoxyglucose F-18 Injection administration.

6 ADVERSE REACTIONS

Hypersensitivity reactions with pruritus, edema and rash have been reported in the postmarketing setting. Have emergency resuscitation equipment and personnel immediately available.

7 DRUG INTERACTIONS

The possibility of interactions of Fludeoxyglucose F-18 Injection with other drugs taken by patients undergoing PET imaging has not been studied.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C

Animal reproduction studies have not been conducted with Fludeoxyglucose F-18 Injection. It is also not known whether Fludeoxyglucose F-18 Injection can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Consider alternative diagnostic tests in a pregnant woman; administer Fludeoxyglucose F-18 Injection only if clearly needed.

8.3 Nursing Mothers

It is not known whether Fludeoxyglucose F-18 Injection is excreted in human milk. Consider alternative diagnostic tests in women who are breast-feeding. Use alternatives to breast feeding (e.g., stored breast milk or infant formula) for at least 10 half-lives of radioactive decay, if Fludeoxyglucose F-18 Injection is administered to a woman who is breast-feeding.

8.4 Pediatric Use


The safety and effectiveness of Fludeoxyglucose F-18 Injection in pediatric patients with epilepsy is established on the basis of studies in adult and pediatric patients. In pediatric patients with epilepsy, the recommended dose is 2.6 mCi. The optimal dose adjustment on the basis of body size or weight has not been determined. In the oncology or cardiology settings, the safety and effectiveness of Fludeoxyglucose F-18 Injection have not been established in pediatric patients.

11 DESCRIPTION

11.1 Chemical Characteristics

Fludeoxyglucose F-18 Injection, USP is a positron emitting radiopharmaceutical that is used for diagnostic purposes in conjunction with positron emission tomography (PET) imaging. The active ingredient 2-deoxy-2- ^{18}F fluoro-D-glucose has the molecular formula of $\text{C}_6\text{H}_{11}^{18}\text{FO}_5$

with a molecular weight of 181.26, and has the following chemical structure:



Fludeoxyglucose F-18 Injection, USP is provided as a ready to use sterile, pyrogen free, clear, colorless citrate buffered solution. Each mL contains between 0.740 to 11.1GBq (20.0-300 mCi) of 2-deoxy-2- ^{18}F fluoro-D-glucose at the EOS, 4.5 mg of sodium chloride in citrate buffer. The pH of the solution is between 4.5 and 7.5. The solution is packaged in a multiple-dose glass vial and does not contain any preservative.

11.2 Physical Characteristics

Fluorine F-18 decays by emitting positron to Oxygen O 18 (stable) and has a physical half-life of 109.7 minutes. The principal photons useful for imaging are the dual 511 keV gamma photons, that are produced and emitted simultaneously in opposite direction when the positron interacts with an electron (Table 2).

Table 2. Principal Radiation Emission Data for Fluorine F-18

Radiation/Emission	% Per Disintegration	Mean Energy
--------------------	----------------------	-------------

Positron(β^+)	96.73	249.8 keV
Gamma(\pm)*	193.46	511.0 keV

*Produced by positron annihilation

From: Kocher, D.C. Radioactive Decay Tables DOE/TIC-I 1026, 89 (1981)

The specific gamma ray constant (point source air kerma coefficient) for fluorine F-18 is 5.7 R/hr/mCi (1.35×10^{-6} Gy/hr/kBq) at 1 cm. The half-value layer (HVL) for the 511 keV photons is 4 mm lead (Pb). The range of attenuation coefficients for this radionuclide as a function of lead shield thickness is shown in Table 3. For example, the interposition of an 8 mm thickness of Pb, with a coefficient of attenuation of 0.25, will decrease the external radiation by 75%.

Table 3. Radiation Attenuation of 511 keV Photons by lead (Pb) shielding

Shield thickness (Pb) mm	Coefficient of attenuation
0	0.00
4	0.50
8	0.25
13	0.10
26	0.01
39	0.001
52	0.0001

For use in correcting for physical decay of this radionuclide, the fractions remaining at selected intervals after calibration are shown in Table 4.

Table 4. Physical Decay Chart for Fluorine F-18

Minutes	Fraction Remaining
0*	1.000
15	0.909
30	0.826
60	0.683
110	0.500
220	0.250

*calibration time

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Fludeoxyglucose F-18 is a glucose analog that concentrates in cells that rely upon glucose as an energy source, or in cells whose dependence on glucose increases under pathophysiological conditions. Fludeoxyglucose F-18 is transported through the cell membrane by facilitative glucose transporter proteins and is phosphorylated within the cell to [^{18}F] FDG-6-phosphate by the enzyme hexokinase. Once phosphorylated it cannot exit until it is dephosphorylated by glucose-6-phosphatase. Therefore, within a given tissue or pathophysiological process, the retention and clearance of Fludeoxyglucose F-

18 reflect a balance involving glucose transporter, hexokinase and glucose-6-phosphatase activities. When allowance is made for the kinetic differences between glucose and Fludeoxyglucose F-18 transport and phosphorylation (expressed as the "lumped constant" ratio), Fludeoxyglucose F-18 is used to assess glucose metabolism.

In comparison to background activity of the specific organ or tissue type, regions of decreased or absent uptake of Fludeoxyglucose F-18 reflect the decrease or absence of glucose metabolism. Regions of increased uptake of Fludeoxyglucose F-18 reflect greater than normal rates of glucose metabolism.

12.2 Pharmacodynamics

Fludeoxyglucose F-18 Injection is rapidly distributed to all organs of the body after intravenous administration. After background clearance of Fludeoxyglucose F-18 Injection, optimal PET imaging is generally achieved between 30 to 40 minutes after administration.

In cancer, the cells are generally characterized by enhanced glucose metabolism partially due to (1) an increase in activity of glucose transporters, (2) an increased rate of phosphorylation activity, (3) a reduction of phosphatase activity or, (4) a dynamic alteration in the balance among all these processes. However, glucose metabolism of cancer as reflected by Fludeoxyglucose F-18 accumulation shows considerable variability. Depending on tumor type, stage, and location, Fludeoxyglucose F-18 accumulation may be increased, normal, or decreased. Also, inflammatory cells can have the same variability of uptake of Fludeoxyglucose F-18.

In the heart, under normal aerobic conditions, the myocardium meets the bulk of its energy requirements by oxidizing free fatty acids. Most of the exogenous glucose taken up by the myocyte is converted into glycogen. However, under ischemic conditions, the oxidation of free fatty acids decreases, exogenous glucose becomes the preferred myocardial substrate, glycolysis is stimulated, and glucose taken up by the myocyte is metabolized immediately instead of being converted into glycogen. Under these conditions, phosphorylated Fludeoxyglucose F-18 accumulates in the myocyte and can be detected with PET imaging.

In the brain, cells normally rely on aerobic metabolism. In epilepsy, the glucose metabolism varies. Generally, during a seizure, glucose metabolism increases. Interictally, the seizure focus tends to be hypometabolic.

12.3 Pharmacokinetics

Distribution: In four healthy male volunteers, receiving an intravenous administration of 30 seconds in duration, the arterial blood level profile for Fludeoxyglucose F-18 decayed triexponentially. The effective half-life ranges of the three phases were 0.2-0.3 minutes, 10-13 minutes with a mean and standard deviation (STD) of 11.6 (\pm) 1.1 min, and 80-95 minutes with a mean and STD of 88 (\pm) 4 min.

Plasma protein binding of Fludeoxyglucose F-18 has not been studied.

Metabolism: Fludeoxyglucose F-18 is transported into cells and phosphorylated to [^{18}F]-FDG-6-phosphate at a rate proportional to the rate of glucose utilization within that tissue. [^{18}F]-FDG-6-phosphate presumably is metabolized to 2-deoxy-2- [^{18}F]fluoro-6-phospho-Dmannose ([^{18}F]FDM-6-phosphate).

Fludeoxyglucose F-18 Injection may contain several impurities (e.g., 2-deoxy-2-chloro-Dglucose (CIDG)). Biodistribution and metabolism of CIDG are presumed to be similar to Fludeoxyglucose F-18 and would be expected to result in intracellular formation of 2-deoxy-2-chloro-6-phospho-D-glucose (CIDG-6-phosphate) and 2-deoxy-2-chloro-6-phospho-D-mannose (CIDM-6-phosphate). The phosphorylated deoxyglucose compounds are dephosphorylated and the resulting compounds (FDG, FDM, CIDG, and CIDM) presumably leave cells by passive diffusion. Fludeoxyglucose F-18 and related compounds are cleared from non-cardiac tissues within 3 to 24 hours after administration. Clearance from the cardiac tissue may require more than 96 hours. Fludeoxyglucose F-18 that is not

involved in glucose metabolism in any tissue is then excreted in the urine.

Elimination: Fludeoxyglucose F-18 is cleared from most tissues within 24 hours and can be eliminated from the body unchanged in the urine. Three elimination phases have been identified in the reviewed literature. Within 33 minutes, a mean of 3.9% of the administered radioactive dose was measured in the urine. The amount of radiation exposure of the urinary bladder at two hours post-administration suggests that 20.6% (mean) of the radioactive dose was present in the bladder.

Special Populations:

The pharmacokinetics of Fludeoxyglucose F-18 Injection have not been studied in renally-impaired, hepatically impaired or pediatric patients. Fludeoxyglucose F-18 is eliminated through the renal system. Avoid excessive radiation exposure to this organ system and adjacent tissues.

The effects of fasting, varying blood sugar levels, conditions of glucose intolerance, and diabetes mellitus on Fludeoxyglucose F-18 distribution in humans have not been ascertained [see *Warnings and Precautions (5.2)*].

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Animal studies have not been performed to evaluate the Fludeoxyglucose F-18 Injection carcinogenic potential, mutagenic potential or effects on fertility.

14 CLINICAL STUDIES

14.1 Oncology

The efficacy of Fludeoxyglucose F-18 Injection in positron emission tomography cancer imaging was demonstrated in 16 independent studies. These studies prospectively evaluated the use of Fludeoxyglucose F-18 in patients with suspected or known malignancies, including non-small cell lung cancer, colo-rectal, pancreatic, breast, thyroid, melanoma, Hodgkin's and non-Hodgkin's lymphoma, and various types of metastatic cancers to lung, liver, bone, and axillary nodes. All these studies had at least 50 patients and used pathology as a standard of truth. The Fludeoxyglucose F-18 Injection doses in the studies ranged from 200 MBq to 740 MBq with a median and mean dose of 370 MBq.

In the studies, the diagnostic performance of Fludeoxyglucose F-18 Injection varied with the type of cancer, size of cancer, and other clinical conditions. False negative and false positive scans were observed. Negative Fludeoxyglucose F-18 Injection PET scans do not exclude the diagnosis of cancer. Positive Fludeoxyglucose F-18 Injection PET scans can not replace pathology to establish a diagnosis of cancer. Non-malignant conditions such as fungal infections, inflammatory processes and benign tumors have patterns of increased glucose metabolism that may give rise to false-positive scans. The efficacy of Fludeoxyglucose F-18 Injection PET imaging in cancer screening was not studied.

14.2 Cardiology

The efficacy of Fludeoxyglucose F-18 Injection for cardiac use was demonstrated in ten independent, prospective studies of patients with coronary artery disease and chronic left ventricular systolic dysfunction who were scheduled to undergo coronary revascularization. Before revascularization, patients underwent PET imaging with Fludeoxyglucose F-18 Injection (74 – 370 MBq, 2 – 10 mCi) and perfusion imaging with other diagnostic radiopharmaceuticals. Doses of Fludeoxyglucose F-18 Injection ranged from 74-370 MBq (2-10 mCi). Segmental, left ventricular, wall-motion assessments of asynergic areas made before revascularization were compared in a blinded manner to assessments made after successful revascularization to identify myocardial segments with functional recovery.

Left ventricular myocardial segments were predicted to have reversible loss of systolic function if they

showed Fludeoxyglucose F-18 accumulation and reduced perfusion (i.e., flow-metabolism mismatch). Conversely, myocardial segments were predicted to have irreversible loss of systolic function if they showed reductions in both Fludeoxyglucose F-18 accumulation and perfusion (i.e., matched defects).

Findings of flow-metabolism mismatch in a myocardial segment may suggest that successful revascularization will restore myocardial function in that segment. However, false-positive tests occur regularly, and the decision to have a patient undergo revascularization should not be based on PET findings alone. Similarly, findings of a matched defect in a myocardial segment may suggest that myocardial function will not recover in that segment, even if it is successfully revascularized. However, false-negative tests occur regularly, and the decision to recommend against coronary revascularization, or to recommend a cardiac transplant, should not be based on PET findings alone. The reversibility of segmental dysfunction as predicted with Fludeoxyglucose F-18 PET imaging depends on successful coronary revascularization. Therefore, in patients with a low likelihood of successful revascularization, the diagnostic usefulness of PET imaging with Fludeoxyglucose F-18 Injection is more limited.

14.3 Neurology

In a prospective, open label trial, Fludeoxyglucose F-18 Injection was evaluated in 86 patients with epilepsy. Each patient received a dose of Fludeoxyglucose F-18 Injection in the range of 185-370 MBq (5-10 mCi). The mean age was 16.4 years (range: 4 months - 58 years; of these, 42 patients were less than 12 years and 16 patients were less than 2 years old). Patients had a known diagnosis of complex partial epilepsy and were under evaluation for surgical treatment of their seizure disorder. Seizure foci had been previously identified on ictal EEGs and sphenoidal EEGs. Fludeoxyglucose F-18 Injection PET imaging confirmed previous diagnostic findings in 16% (14/87) of the patients; in 34% (30/87) of the patients, Fludeoxyglucose F-18 Injection PET images provided new findings. In 32% (27/87), imaging with Fludeoxyglucose F-18 Injection was inconclusive. The impact of these imaging findings on clinical outcomes is not known.

Several other studies comparing imaging with Fludeoxyglucose F-18 Injection results to subsphenoidal EEG, MRI and/or surgical findings supported the concept that the degree of hypometabolism corresponds to areas of confirmed epileptogenic foci. The safety and effectiveness of Fludeoxyglucose F-18 Injection to distinguish idiopathic epileptogenic foci from tumors or other brain lesions that may cause seizures have not been established.

15 REFERENCES

1. Gallagher B.M., Ansari A., Atkins H., Casella V., Christman D.R., Fowler J.S., Ido T., MacGregor R.R., Som P., Wan C.N., Wolf A.P., Kuhl D.E., and Reivich M. "Radiopharmaceuticals XXVII. 18F-labeled 2-deoxy-2-fluoro-D-glucose as a radiopharmaceutical for measuring regional myocardial glucose metabolism in vivo: tissue distribution and imaging studies in animals," *J Nucl Med*, 1977; 18, 990-6.
2. Jones S.C., Alavi, A., Christman D., Montanez, I., Wolf, A.P., and Reivich M. "The radiation dosimetry of 2 [F-18] fluoro-2-deoxy-D-glucose in man," *J Nucl Med*, 1982; 23, 613-617.
3. Kocher, D.C. "Radioactive Decay Tables: A handbook of decay data for application to radiation dosimetry and radiological assessments," 1981, DOE/TIC-I 1026, 89.
4. ICRP Publication 53, Volume 18, No. 1-4, 1987, pages 75-76.

16 HOW SUPPLIED/STORAGE AND DRUG HANDLING

Fludeoxyglucose F-18 Injection, USP is supplied in a multi-dose, capped 50 mL glass vial containing between 0.740 – 11.1 GBq/mL (20 - 300 mCi/mL), of no carrier added 2-deoxy-2-[18F] fluoro-D-glucose, at end of synthesis, in approximately 23 mL or 35 mL. The contents of each vial are sterile, pyrogen-free and preservative-free.

NDC 67939-010-35

NDC 67939-010-23

Store the Fludeoxyglucose F-18 Injection, USP vial upright in a lead shielded container at 25°C (77°F); excursions permitted to 15-30°C (59-86°F).

Store and dispose of Fludeoxyglucose F-18 Injection, USP in accordance with the regulations and a general license, or its equivalent, of an Agreement State or a Licensing State.

The expiration date and time are provided on the container label. Use Fludeoxyglucose F18 Injection, USP within 12 hours from the EOS time.

17 PATIENT COUNSELING INFORMATION

Instruct patients in procedures that increase renal clearance of radioactivity. Encourage patients to:

- drink water or other fluids (as tolerated) in the 4 hours before their PET study.
- void as soon as the imaging study is completed and as often as possible thereafter for at least one hour.

Manufactured by:	The University of Utah DBA Cyclotron Radiochemistry Lab Hunstman Cancer Institute 2000 Circle of Hope Salt Lake City, UT 84112
Distributed by:	The University of Utah DBA Cyclotron Radiochemistry Lab Hunstman Cancer Institute 2000 Circle of Hope Salt Lake City, UT 84112

Outer Package Labels & Vial Labels

Fludeoxyglucose F-18 Injection 35 mL (67939-010-35)

NDC# 67939-010-35

Multiple-Dose Vial

Fludeoxyglucose F 18 Injection, USP

20 mCi/mL to 300 mCi/mL @ EOS*

Activity @ EOS*: Total _____ mCi Volume: _____ mL
Concentration: _____ mCi/mL

Sterile, Non-pyrogenic

Calibration (EOS*) Time _____
Calibration Date: _____

Diagnostic - For Intravenous Use Only

Exp. Date/Time _____, _____
LOT# _____

(Expires 12 hours after EOS*)

Contains:

Each ml contains 0.74 GBq to 11.1 GBq
(20 mCi to 300 mCi) of no-carrier added
Fludeoxyglucose F 18 and 4.5 mg of sodium chloride
in citrate buffer.

Store at 25°C (77°F) (see insert)
Store upright in a shielded container.
Aseptically withdraw and handle doses.
[18F] Half-Life = 109.7 minutes
Calculate correct dosage from date and
time of calibration.

Do not use if cloudy or if it contains particulate matter

*EOS = End of Synthesis

CAUTION: RADIOACTIVE MATERIAL 

Manufactured by: The University of Utah
DBA Cyclotron Radiochemistry Lab
Huntsman Cancer Institute
Salt Lake City, UT, 84112

Rx ONLY

NDC# 67939-010-35

Multiple-Dose Vial

Fludeoxyglucose F 18 Injection, USP

20 mCi/mL to 300 mCi/mL @ EOS*

Activity @ EOS*: Total _____ mCi Volume: _____ mL
Concentration: _____ mCi/mL

Sterile, Non-pyrogenic

Calibration (EOS*) Time _____
Calibration Date: _____

Diagnostic - For Intravenous Use Only

Exp. Date/Time _____, _____
LOT# _____

(Expires 12 hours after EOS*)

Contains:

Each ml contains 0.74 GBq to 11.1 GBq
(20 mCi to 300 mCi) of no-carrier added
Fludeoxyglucose F 18 and 4.5 mg of sodium chloride
in citrate buffer.

Store at 25°C (77°F) (see insert)
Store upright in a shielded container.
Aseptically withdraw and handle doses.
[18F] Half-Life = 109.7 minutes
Calculate correct dosage from date and
time of calibration.

Do not use if cloudy or if it contains particulate matter

*EOS = End of Synthesis

CAUTION: RADIOACTIVE MATERIAL 

Manufactured by: The University of Utah
DBA Cyclotron Radiochemistry Lab
Huntsman Cancer Institute
Salt Lake City, UT, 84112

Rx ONLY

Fludeoxyglucose F-18 Injection 23 mL (67939-010-23)

NDC# 67939-010-23

Multiple-Dose Vial

Fludeoxyglucose F 18 Injection, USP

20 mCi/mL to 300 mCi/mL @ EOS*

Activity @ EOS*: Total _____ mCi Volume: _____ mL
Concentration: _____ mCi/mL

Sterile, Non-pyrogenic

Calibration (EOS*) Time _____

Calibration Date: _____

Diagnostic - For Intravenous Use Only

Exp. Date/Time _____, _____

LOT# _____

(Expires 12 hours after EOS*)

Contains:

Each ml contains 0.74 GBq to 11.1 GBq
(20 mCi to 300 mCi) of no-carrier added
Fludeoxyglucose F 18 and 4.5 mg of sodium chloride
in citrate buffer.

Store at 25°C (77°F) (see insert)
Store upright in a shielded container.
Aseptically withdraw and handle doses.
[18F] Half-Life = 109.7 minutes
Calculate correct dosage from date and
time of calibration.

Do not use if cloudy or if it contains particulate matter
*EOS = End of Synthesis

CAUTION: RADIOACTIVE MATERIAL



Manufactured by: The University of Utah
DBA Cyclotron Radiochemistry Lab
Huntsman Cancer Institute
Salt Lake City, UT, 84112

Rx ONLY

NDC# 67939-010-23

Multiple-Dose Vial

Fludeoxyglucose F 18 Injection, USP

20 mCi/mL to 300 mCi/mL @ EOS*

Activity @ EOS*: Total _____ mCi Volume: _____ mL
Concentration: _____ mCi/mL

Sterile, Non-pyrogenic

Calibration (EOS*) Time _____

Calibration Date: _____

Diagnostic - For Intravenous Use Only

Exp. Date/Time _____, _____

LOT# _____

(Expires 12 hours after EOS*)

Contains:

Each ml contains 0.74 GBq to 11.1 GBq
(20 mCi to 300 mCi) of no-carrier added
Fludeoxyglucose F 18 and 4.5 mg of sodium chloride
in citrate buffer.

Store at 25°C (77°F) (see insert)
Store upright in a shielded container.
Aseptically withdraw and handle doses.
[18F] Half-Life = 109.7 minutes
Calculate correct dosage from date and
time of calibration.

Do not use if cloudy or if it contains particulate matter
*EOS = End of Synthesis

CAUTION: RADIOACTIVE MATERIAL



Manufactured by: The University of Utah
DBA Cyclotron Radiochemistry Lab
Huntsman Cancer Institute
Salt Lake City, UT, 84112

Rx ONLY

FLUDEOXYGLUCOSE F 18

fludeoxyglucose f 18 injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:67939-010
Route of Administration	INTRAVENOUS	DEA Schedule	

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FLUDEOXYGLUCOSE F-18 (UNII: 0Z5B2CJX4D) (FLUDEOXYGLUCOSE F-18 - UNII:0Z5B2CJX4D)	FLUDEOXYGLUCOSE F-18	300 mCi in 1 mL

Inactive Ingredients

Ingredient Name	Strength
SODIUM CHLORIDE (UNII: 451W47IQ8X)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67939-010-35	35 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		
2	NDC:67939-010-23	23 mL in 1 VIAL, MULTI-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA204498	11/01/2013	

Labeler - The University of Utah DBA Cyclotron Radiochemistry Lab / Huntsman Cancer Institute (018432646)

Establishment

Name	Address	ID/FEI	Business Operations
The University of Utah DBA Cyclotron Radiochemistry Lab / Huntsman Cancer Institute		018432646	positron emission tomography drug production(67939-010) , manufacture(67939-010)

Revised: 6/2015

The University of Utah DBA Cyclotron Radiochemistry Lab / Huntsman Cancer Institute